



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(19)

(11) Publication number:

**0 204 480**  
**A2**

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: **86303978.0**

(51) Int. Cl.<sup>4</sup>: **C 07 K 7/00**  
**C 12 P 21/02, C 12 N 15/00**  
**A 61 K 37/02**

(22) Date of filing: **27.05.86**

(30) Priority: **03.06.85 US 740780**

(43) Date of publication of application:  
**10.12.86 Bulletin 86/50**

(84) Designated Contracting States:  
**BE CH DE FR GB IT LI LU NL SE**

(71) Applicant: **ELI LILLY AND COMPANY**  
**Lilly Corporate Center**  
**Indianapolis Indiana 46285(US)**

(72) Inventor: **Dimarchi, Richard Dennis**  
**3815 Wolf Creek Circle**  
**Carmel Indiana 46032(US)**

(72) Inventor: **Brooke, Gerald Stephen**  
**2700 Dellzell Drive**  
**Indianapolis Indiana 46220(US)**

(74) Representative: **Hudson, Christopher Mark et al,**  
**Eri Wood Manor**  
**Windlesham Surrey GU20 6PH(GB)**

(54) **Synthetic vaccines for foot and mouth disease.**

(57) Compounds having activity in treating or preventing foot and mouth disease are disclosed. The compounds contain a sequence of the formula

**X-A-Y-B-Z**

in which A and B are amino acid residue sequences comprising sequences of the foot-and-mouth disease virus VP<sub>1</sub> capsid protein serotypes, one of which contains from 18 to 24 amino acid residues and includes the amino acid residue sequence in positions 141 to 158 of the O serotype or the equivalent sequence of other serotypes and the other of which contains from 14 to 20 amino acid residues and includes the amino acid residue sequence in positions 200 to 213 of the O serotype or the equivalent sequence of other serotypes;

X is H, H-Cys, or H-Cys-Cys, the sulfhydryls of which may be blocked or free;

Z is OH, Cys-OH, or Pro-Cys-Gly-OH, the sulfhydryls of which may be blocked or free; and

Y is a sequence of from about 2 to about 6 amino acid residues, or a pharmaceutically-acceptable salt thereof.

**EP 0 204 480 A2**

## SYNTHETIC VACCINES FOR FOOT AND MOUTH DISEASE

Foot-and-mouth disease (FMD) is a highly contagious, debilitating disease that afflicts cloven-hoofed animals. The causative agent, foot-and-mouth disease virus (FMDV), is a small animal virus of the picornavirus family having a single stranded positive sense RNA genome of about 8000 nucleotides.

In the past, vaccines have been produced for FMD using either an inactivated virus or a live attenuated virus. These approaches, although generally effective, have not been problem-free. Occasions on which the virus was incompletely inactivated or insufficiently attenuated can and have given rise to FMD outbreaks. A synthetic FMDV vaccine which would eliminate all aspects of virus handling in production and vaccination is perceived as a most desirable alternative [D. Rowlands, Endeavor 8 (3), 123-127 (1984)].

Studies of FMDV over the last several years have established that the picornavirus contains primarily four capsid proteins designated VP<sub>1</sub>, VP<sub>2</sub>, VP<sub>3</sub>, and VP<sub>4</sub>. The actual labelling of these VP proteins has been subject to a measure of inconsistency. The relevant protein insofar as this invention is concerned has been designated VP<sub>3</sub> by some research groups [see, e.g., Kleid et al., Science 214, 1125 (1981)] while being labelled VP<sub>1</sub> by others [see, e.g., Clarke et al., FEBS Letters 157, 261 (1983)]. For present purposes this protein is designated VP<sub>1</sub>.

VP<sub>1</sub> has been the subject of considerable investigation and structural analysis, it having been found to elicit a neutralizing antibody response in swine [J. Laporte, J. Grosclaude, J. Wantyghem, S. Bernard, P. Rouze, C. R. Acad. Sci. 276, 3399 (1973)] and to protect both swine and cattle from FMDV infection [H.L. Bachrach, D.M. Moore, P.D. McKercher, J. Polatnick, J. Immunol. 115, 1636 (1975)]. A biosynthetic fusion polypeptide of TrpLE and VP<sub>1</sub>, upon multiple vaccination, has been reported to protect cattle and swine against a viral contact challenge [D.G. Kleid, D. Yansura, B. Small, D. Dowbenko, D. Moore, M. Grubman, P. McKercher D.O. Morgan, E.V. Robertson, H.L. Bachrach, Science 214, 1125-1129 (1981)].

Strohmaier et al., J. Gen. Virol. 59, 295-306 (1982), having chemically and enzymatically digested VP<sub>1</sub>, have concluded that major immunogenic regions capable of raising neutralizing antibodies are present at amino acid sequences 146-154 and 201-213.

Bittle et al., Nature 298, 30-33 (1982) chemically synthesized peptides corresponding to regions 141-160 and 200-213 of the VP<sub>1</sub> polypeptide of FMDV and demonstrated the ability of these peptides to produce high levels of neutralizing antibody in guinea pigs and rabbits. A report on this work is contained also in the PCT International Patent Application No. WO 83 03,547, published October 27, 1983.

A major theme throughout all of the foregoing work with the small peptide sequences is the belief and finding that their immunogenic effect, if present at all,

is achieved only upon the linking of the peptide to a high molecular weight carrier, most often keyhole limpet hemocyanin (KLH) carrier.

PCT Application No. WO 83 03,547 suggests that  
5 a small peptide, in monomeric and uncyclized form, is incapable of eliciting a neutralizing index sufficient to achieve viral protection. Specifically, the publication states that a neutralizing index of about 1.5 or  
10 greater is required to protect an animal against the virus and that the neutralizing index of the monomeric, unconjugated peptide having a sequence which corresponds substantially to the amino acid residue sequence in positions about 141 to about 160 of O<sub>1</sub>K FMDV is approximately 0.5. This result is to be compared with neu-  
15 tralizing indices of about 3.7 to about 3.9 reported for the same peptide when conjugated to KLH.

A class of novel, monomeric, unconjugated, small peptides has now been discovered. These peptides, administered free of any carrier, are capable of eliciting an unforeseen markedly enhanced neutralization re-  
20 sponse against FMDV. These peptides have induced unprecedented protection in cattle with single vaccination and complete protection after repeat immunization. Until the present discovery, this latter achievement has  
25 been proven only with antigens approximately five times larger in size than these newly discovered peptides.

In accordance with the invention, there is provided a compound containing a sequence of Formula (I):

## X-A-Y-B-Z (I)

in which A and B are amino acid residue sequences comprising sequences of the foot-and-mouth disease virus VP<sub>1</sub> capsid protein serotypes, one of which contains from 18 to 24 amino acid residues and includes the amino acid residue sequence in positions 141 to 158 of the O serotype or the equivalent sequence of other serotypes and the other of which contains from 14 to 20 amino acid residues and includes the amino acid residue sequence in positions 200 to 213 of the O serotype or the equivalent sequence of other serotypes;

X is H, H-Cys, or H-Cys-Cys, the sulfhydryls of which may be blocked or free;

Z is OH, Cys-OH, or Pro-Cys-Gly-OH, the sulfhydryls of which may be blocked or free; and

Y is a sequence of from about 2 to about 6 amino acid residues, or a pharmaceutically-acceptable salt thereof.

As noted, this invention is directed to a class of synthetic, immunogenic peptides of Formula (I):

## X-A-Y-B-Z. (I)

The groups A and B include amino acid sequences corresponding to selected sequences present in the VP<sub>1</sub> capsid protein of FMDV serotype O, subtype 1, strain Kaufbeuren (O<sub>1</sub>K) and equivalent analogous sequences present in any of other foot-and-mouth disease virus serotypes.

Seven serotypes of FMDV are known to exist. The most common are European and carry designations A, O, and C. Others are three South African, SAT-1, SAT-2, and SAT-3, and one Asian, Asia-1. Each serotype in addition is known to have a number of subtypes and associated subtype strains. Specific examples are serotype O, subtype 1, strain Kaufbeuren (O<sub>1</sub>K); serotype O, subtype 1, strain Campos (O<sub>1</sub>C); serotype O, subtype 1, strain British Field Strain (O<sub>1</sub>BFS); serotype A, subtype 10, strain 61 (A<sub>10</sub>,61); serotype A, subtype 12, strain 119 (A<sub>12</sub>,119); serotype A, subtype 24, strain Cruzeiro (A<sub>24</sub>C); serotype A, subtype 27 (A<sub>27</sub>); serotype A, subtype 79 (A<sub>79</sub>); and serotype C, subtype 3, strain Indaial (C<sub>3</sub>I).

The amino acid residue sequences in positions 141-158 and 200-213 of the O serotype represent the focus of the definitions of groups A and B of the peptides of this invention. In any specific instance group A includes one of the sequences and group B the other. In addition, the analogous but not necessarily identical sequences present in any of the other serotype and serotype subtypes are included within the definition of the groups A and B. In those cases in which the definition of the groups A and B includes a length of amino acid residues greater than 18 and 14 amino acids for the sequences corresponding to amino acids 141-158 and 200-213 of the O serotype, respectively, the remaining residues can be any amino acid.

For purposes of illustration, the following are provided as examples of the sequence 141-158 for O<sub>1</sub>K

X-6060

-6-

and analogous sequences of other viral serotypes and subtypes:

	141	150	158
O <sub>1</sub> K	ValProAsnLeuArgGlyAspLeuGlnValLeuAlaGlnLysValAlaArgThr		
5 O <sub>1</sub> BFS	ValProAsnLeuArgGlyAspLeuGlnValLeuAlaGlnLysValAlaArgThr		
O <sub>1</sub> C	ValProAsnValArgGlyAspLeuGlnValLeuAlaGlnLysValAlaArgThr		
A10,61	Ser - ArgSer - GlyAspLeuGlySerIleAlaAlaArgValAlaThrGln		
A12,119	SerGly - ValArgGlyAspPheGlySerLeuAlaProArgValAlaArgGln		
A24C	SerGly - ArgArgGlyAspMetGlyThrLeuAlaAlaArgValValLysGln		
10 A27	Gln - - ArgAlaGlyAspMetGlySerLeuAlaAlaArgValAlaLysGln		
A79	SerGly - ArgArgGlyAspMetGlySerLeuAlaAlaArgValAlaLysGln		
C3I	- - - ArgArgGlyAspLeuValHisLeuAlaAlaAlaHisAlaArgHis		

The following are examples of the sequence  
 15 200-213 for O<sub>1</sub>K and analogous sequences of other viral serotypes and subtypes:

	200	213
O <sub>1</sub> K	ArgHisLysGlnLysIleValAlaProValLysGlnThrLeu	
20 O <sub>1</sub> BFS	ArgHisLysGlnLysIleValAlaProValLysGlnThrLeu	
A10,61	ArgTyrLysGlnLysIleIleAlaProAlaLysGlnLeuLeu	
A12,119	ArgHisLysGlnLysIleIleAlaProGlyLysGln - Leu	
A24C	ArgHisLysGlnLysIleIleAlaProAlaLysGlnLeuLeu	
C3I	ArgHisLysGlnProLeuIleAlaProAlaLysGlnLeuLeu	
25		

In the peptides provided by this invention,  
 one of the groups A or B contains a peptide sequence  
 141-158 and the other a peptide sequence 200-213.  
 Preferably, one of the groups A or B is a peptide  
 30 sequence 141-158 and the other is a peptide sequence

200-213. More preferably, group A is the sequence 200-213 and group B the sequence 141-158. Even more preferably, the groups A and B of the peptides of this invention are peptide sequences derived from the same FMDV serotype.

The group X represents the amino terminal portion of the peptides of this invention. X can represent the hydrogen of the amino terminus or a mono-cysteinyl or di-cysteinyl extension. If X is mono- or di-cysteinyl, the cysteine sulfhydryls may be blocked or free. Any of a wide range of blocking groups familiar to those skilled in the art can be used. Examples of these are sulfonate, carboxymethyl, carboxamidomethyl, aminoethyl, and the like. Preferably, however, X is hydrogen.

The group Z defines the carboxyl terminal of the peptides of this invention. Z thus is a hydroxyl or a cysteine-containing extension, viz., cysteine or the tripeptide Pro-Cys-Gly-OH. Again, the cysteine may be blocked or free, and, if blocked, can be so blocked using any of the same blocking groups previously described.

The group Y links groups A and B and represents an amino acid residue or a sequence having from 2 to 6 amino acid residues. Preferably, Y is an amino acid sequence having from 2 to 4 amino acid residues. The sequence can be comprised of any of a wide range of amino acid residues, proline being an amino acid residue of choice, and sequences containing Pro-Pro being particularly preferred. Specific preferred sequences may be Pro-Pro-Ser, or Leu-Pro-Pro-Thr.



Included in the compounds of this invention are their pharmaceutically acceptable acid addition salts and their pharmaceutically acceptable carboxylic acid salts. "Pharmaceutically acceptable salts" refers to those salts useful in the chemotherapy of a warm-blooded animal.

The term "pharmaceutically acceptable acid addition salts" encompasses both organic and inorganic acid addition salts including, for example, those prepared from acids such as hydrochloric, hydrofluoric, sulfuric, sulfonic, tartaric, fumaric, hydrobromic, glycolic, citric, maleic, phosphoric, succinic, acetic, nitric, benzoic, ascorbic, p-toluenesulfonic, benzenesulfonic, naphthalenesulfonic, propionic, and the like. Preferably, the acid addition salts are those prepared from hydrochloric acid, acetic acid, or succinic acid. The above salts are prepared by conventional methods known to those skilled in the art.

The term "carboxylic acid salts" includes, for example, ammonium, alkali metal salts such as sodium, potassium, and lithium, and the like.

For convenience, the amino acids referred to are described by their approved shorthand three-letter designations.

These designations are as follows:

3-letter

Alanine	Ala
Arginine	Arg
Asparagine	Asn
Aspartic Acid	Asp

		<u>3-letter</u>
	Cysteine	Cys
	Glutamic Acid	Glu
	Glutamine	Gln
5	Glycine	Gly
	Histidine	His
	Isoleucine	Ile
	Leucine	Leu
	Lysine	Lys
10	Phenylalanine	Phe
	Proline	Pro
	Serine	Ser
	Threonine	Thr
	Tryptophan	Trp
15	Tyrosine	Tyr
	Valine	Val

Examples of compounds of this invention are:

	VP <sub>1</sub> (O <sub>1</sub> K)	H[(200-213)ProProSer(141-158)]ProCysGly-OH;
	VP <sub>1</sub> (O <sub>1</sub> K)	H-Cys[(200-213)ProProSer(141-158)]Cys-OH;
20	VP <sub>1</sub> (O <sub>1</sub> K)	H-CysCys[(200-213)ProProSer(141-158)]- ProCysGly-OH;
	VP <sub>1</sub> (O <sub>1</sub> K)	H[(200-213)ProProSer(141-158)]OH;
	VP <sub>1</sub> (O <sub>1</sub> BFS)	H[(141-158)ProPro(200-213)]OH;
	VP <sub>1</sub> (A <sub>10,61</sub> )	H[(141-158)LeuProProSer(200-213)]OH;
25	VP <sub>1</sub> (A <sub>12,119</sub> )	H-Cys[(200-213)LeuProProSer(141-158)]OH;
	VP <sub>1</sub> (A <sub>24</sub> C)	H[(200-213)ValProProThrArg(141-158)]OH;
	VP <sub>1</sub> (O <sub>1</sub> C)	H-CysCys[(141-158)ProPro(200-213)]Cys-OH;
	VP <sub>1</sub> (C <sub>3</sub> I)	H[(200-213)ProProSer(141-158)]OH;
	VP <sub>1</sub> (O <sub>1</sub> K)	H-Cys-[ArgTyrAsnArgAsnAla(141-158)]Leu- ProProThr[GluAla(200-213)]OH; and the like.
30		

The compounds of this invention can be prepared by any of a variety of recognized peptide synthesis techniques including classical (solution) methods, solid phase methods, and the more recently available recombinant DNA methods.

One of the principal methods for preparing the compounds of this invention is by the solid phase technique in which the amino acid sequence is constructed sequentially from an initial, insoluble, resin-supported C-terminal amino acid. Techniques for the solid phase method are described by J. Stewart et al., Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, Illinois, 1984.

Specific examples in state of the art solid phase peptide synthesis which may be utilized in the preparation of these FMDV synthetic immunogens are provided in reports by R.D. DiMarchi, J.P. Tam, R.B. Merrifield, Int. J. Pep. Prot. Res. 19, 270-279 (1982); R.B. Merrifield, L.D. Vizioli, H.G. Boman, Biochem. 21, 5020-5031 (1982); S. Mojsov, R.B. Merrifield, Eur. J. Biochem. 145, 601-605 (1984). The more important features are the following:

1. Utilization of copoly(styrene - 1% divinylbenzene) resin, 200-400 mesh beads, available from Bio-Rad laboratories.
2. Functionalization of the resin support to aminomethyl-resin by the method of Mitchell et al., 1978 (see below).
3. Amidation of the carboxy-terminal amino acid to the aminomethyl-resin as its t-butyloxy-

carbonyl-aminoacyl-[4-(oxymethyl)phenyl]acetic acid derivative.

4. Cleavage of the amino-protecting t-butyloxy-carbonyl (Boc) group in 50% TFA/CH<sub>2</sub>Cl<sub>2</sub> and neutralization of the amine in 5% DIEA/CH<sub>2</sub>Cl<sub>2</sub>.
5. Double coupling at each residue is utilized. With the exception of Asn, Gln, and Arg the first coupling is by preformed symmetrical anhydride. The second coupling in all cases is by preformed HOBt-ester. The first coupling for the aforementioned exceptions is analogous to the second coupling.
6. The peptide is cleaved from the resin support by exposure to an acid in the presence of a carbonium ion scavenger.
7. The following side-chain protecting groups preferably are utilized: Arginine (tosyl), Aspartic acid (O-cyclopentyl), Cysteine (4-methoxybenzyl), Glutamic acid (O-cyclopentyl), Histidine (benzyloxymethyl), Lysine (2-chlorobenzyloxycarbonyl), Serine (benzyl), Threonine (benzyl), Tryptophan (formyl) and Tyrosine (2-bromobenzyloxycarbonyl).

The preferred peptidyl-resin cleavage conditions involve exposing the peptidyl-resin to anhydrous HF at a temperature of from -10°C to +15°C. The preferred carbonium ion scavenger is p-cresol.

Compounds of this invention can also be prepared via recombinant DNA methodology. In their preparation, a nucleotide sequence coding for the

desired peptide is prepared using now routine methods for such synthesis. These methods generally involve preparation of oligonucleotides coding both for fragments of the desired coding sequence and for the complementary sequence thereof. The oligonucleotides are designed to provide overlap of one fragment of the coding sequence with two fragments of the complementary sequence and vice versa. The oligonucleotides are paired and joined, ultimately producing the desired gene sequence.

The sequence is inserted into a cloning vector at a location which permits expression of the peptide product for which it codes. A suitable cloning vector contains at least a portion of a gene's expression control sequence.

A typical expression control sequence can be described in terms of five elements. In the order in which they appear in the gene, the elements are as follows: (a) the promoter region; (b) the 5' untranslated region; (c) the protein coding sequence; (d) the 3' untranslated region; and (e) the transcription termination site.

The function of each of these elements in gene systems is well recognized. The promoter region mediates initiation of messenger RNA (mRNA) production (transcription). The promoter may be (1) free of external control (constitutive), (2) under the control of a repressor, a substance that, when present, represses gene function, or (3) under the control of an inducer, a substance that is required to induce gene function.

The lipoprotein (lpp) gene, for example, is free from external control and thus is termed "constitutive".

5 Located at or near the promoter is the "transcription initiation site", a point at which RNA polymerase binds to initiate transcription of mRNA. Once transcription is initiated, mRNA is produced. The structure of the resulting mRNA is determined by the DNA sequences of the gene elements (b) to (d) above.

10 The resulting mRNA carries a sequence which is translatable into protein product. The translatable sequence is located downstream from the 5' untranslated region and upstream from the 3' untranslated region. Translation is mediated by the binding of ribosomes to a sequence in the mRNA 5' untranslated region denoted as  
15 the ribosome binding site and is initiated at the translation start codon (AUG) appearing as the first codon of the product gene sequence and coding as well for the amino acid methionine (Met). Translation terminates at one or more termination codons appearing  
20 at the end of the translation region.

By the techniques of recombinant DNA, it has become possible to prepare cloning vectors useful for the production of selected foreign (exogenous) proteins by inserting into such vectors an expression control  
25 sequence, i.e., a sequence of nucleotides that controls and regulates expression of structural genes with production of exogenous protein when operatively linked to those genes.

30 In the context of the foregoing, the term "expression control sequence" includes elements (a), (b), (d), and (e) above.

Recombinant DNA methodology can be employed to express compounds of this invention either as a portion of a larger "hybrid" molecule or by direct expression. In the direct expression mode, the cloning vector is  
5 designed such that the expression product is composed entirely of desired product preceded by a methionine (Met) residue resulting from the presence of the essential start codon. The superfluous Met residue can be removed by treating the product with cyanogen bromide or  
10 with phenyl isothiocyanate followed by a strong anhydrous acid, such as trifluoroacetic acid.

In the hybrid molecule expression mode, a DNA sequence coding for the desired product is inserted into the expression control sequence of a cloning vector at a  
15 point such that the product expressed comprises a hybrid protein. "Hybrid protein" means a recombinant DNA product comprising a foreign protein, generally all or a portion of the natural (endogenous) protein produced by the expression control sequence (for example, lipoprotein in the lipoprotein gene), to which is attached the  
20 desired protein.

The properly designed hybrid protein produced by recombinant DNA methodology will contain a cleavage site at the junction of the endogenous protein portion and the desired product. The cleavage site permits  
25 generation of mature product by chemical or enzymatic treatment of the hybrid protein product. Highly useful selective cleavage sites comprise a DNA sequence which codes for an amino acid or a sequence of amino acids  
30 which can be cleaved chemically or enzymatically at its C-terminal.

Examples of chemical agents useful for cleaving proteins are cyanogen bromide, BNPS-skatole, hydroxylamine, and the like. Cyanogen bromide cleaves proteins at the C-terminal of a methionine residue.

5 Therefore, the selective cleavage site is a methionine residue itself.

Hydroxylamine cleaves at the C-terminal of the moiety -Asn-Z- in which Z is Gly, Leu, or Ala.

10 BNPS-skatole cleaves at the C-terminal of a tryptophan residue.

Examples of enzymatic agents useful for cleavage are trypsin, papain, pepsin, plasmin, thrombin, enterokinase, and the like. Each effects cleavage at a particular amino acid sequence which it recognizes. For  
15 example, the selective cleavage site that enterokinase recognizes is the amino acid sequence  $-(\text{Asp})_n\text{-Lys-}$  in which  $n$  is an integer from 2 to 4.

A most preferred selective cleavage site, especially for those compounds of this invention that  
20 lack methionine, is a methionine residue. This residue, joined to the N-terminus of the desired product, is readily cleaved by known methods using cyanogen bromide to produce the desired mature product.

In constructing useful cloning vectors,  
25 several elements are required. Two of the required elements are common to all useful cloning vectors. First, the vector must have a DNA segment containing a functional origin of replication (replicon). Plasmids and phage DNA by their very nature contain replicons  
30 facilitating replication in a host cell.



Secondly, the vector must have a DNA segment which conveys to a transformable host cell a property useful for selection of transformed cells from non-transformed cells. Any of a wide range of properties can be used for selection purposes. One of the most commonly used properties is antibiotic resistance, e.g., tetracycline resistance or ampicillin resistance.

The foregoing two elements generally are present in readily available and recognized cloning vectors. Examples of suitable cloning vectors are bacterial plasmids, such as plasmids from E. coli, including pBR322, pMB9, ColE1, pCR1; wider host range plasmids, including RP4; phage DNAs, such as lambda, and the like. Most, if not all, of the above recognized vectors already carry the aforescribed two elements.

A third element is the expression control sequence. Any of a wide range of such control sequences can be used including, for example, those from the lipoprotein gene, the  $\beta$ -galactosidase gene, the tryptophan gene, the  $\beta$ -lactamase gene, phage lambda, and the like.

In producing a suitable cloning vector by insertion of the selected expression control sequence, routine methods are used. Various sites exist within cloning vectors at which cuts can be made using a restriction endonuclease specific for such site. Any of these sites can be selected for insertion of the expression control sequence. As an example, in the well-recognized and documented plasmid pBR322, several suitable restriction sites exist, any of which may be

employed as insertion sites. A PstI site is located within the gene for  $\beta$ -lactamase. Other sites outside of any specific coding region are EcoRI and PvuII. These and other sites are well recognized by those skilled in the art.

Taking advantage of any of these sites or others, insertion of an expression control sequence or the essential portion thereof can be readily accomplished in production of vectors defined by this invention.

A fourth element, of course, is the DNA sequence coding for the desired product. As previously noted, this DNA sequence can be constructed synthetically, e.g., using the recognized phosphotriester method or other well-recognized methods.

Suitable cloning vectors can be used in a wide range of host organisms, for example, gram-negative prokaryotic organisms such as Escherichia coli, Serratia, Pseudomonas, and the like; gram-positive prokaryotic organisms, such as Bacillus, Streptomyces, and the like; and eukaryotic organisms, such as Saccharomyces, and the like. Preferably, the host organism is a gram-negative prokaryotic organism. Of gram-negative prokaryotic organisms, E. coli is especially preferred, for example, E. coli K-12 strains, such as RV308.

Employing well recognized methodology, the appropriately prepared cloning vectors are used to transform suitable host organisms, are amplified in such organisms, and exogenous protein product is expressed using standard fermentation conditions. The exogenous

protein product is isolated by routine methods from the resulting fermentation broth.

Thus, in accordance with the invention, there also is provided a process for preparing a synthetic  
5 foot and mouth disease vaccine containing a sequence of Formula (I):

X-A-Y-B-Z (I),

10 in which X, A, Y, B and Z are as defined previously, which comprises:

- (a) Cleaving a protecting group from a correspond-  
ingly protected compound of Formula (I); or
- 15 (b) Culturing a microorganism transformed by an expression transfer vector containing a DNA sequence coding for a compound of Formula (I), under conditions suitable for expression of the DNA sequence coding for the compound of  
20 Formula (I), and purifying the Formula (I) compound from the lysate or culture medium of said microorganism; and
- (c) If desired, salifying a free compound to form  
25 a corresponding pharmaceutically-acceptable salt.

The compounds of this invention, active in treating foot and mouth disease, can be used in a variety of pharmaceutical compositions and formulations and can be administered by a variety of conventional  
30 routes, such as intranasal, intramuscular, intravenous, subcutaneous, and intraperitoneal.

In administering the compounds of this invention, the pharmaceutical forms suitable for injection include sterile aqueous solutions or dispersions and sterile powders for reconstitution into sterile injectible solutions or dispersions. The carrier can be a solvent or dispersing medium containing, for example, water, ethanol, polyol (for example glycerol, propylene glycol, liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. Proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be ensured by various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol, m-cresol, sorbic acid, and the like. In many cases, it will be desirable to include isotonic agents, for example, sugars, sodium chloride, and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the compounds of this invention in the required amount of the appropriate solvent with various of the other ingredients, as desired.

If desired, and for more effective distribution, the compounds can be incorporated into slow release or targeted delivery systems such as polymer matrices, liposomes, and microspheres.

Doses of the compounds of this invention are administered to the recipient for a period during which prophylactic or therapeutic treatment for foot and mouth disease is desired. The weight of the recipient and  
5 mode of administration will have an influence upon the size of the dose necessary to induce a particular response.

It is especially advantageous to formulate the compounds of this invention in unit dosage form for ease  
10 of administration and uniformity of dosage. "Unit dosage form" refers to a physically discrete unit suited to provide a single dose for the subject to be treated. Each unit contains a predetermined quantity of the compound calculated to produce the desired therapeutic  
15 or prophylactic effect in association with the pharmaceutically acceptable carrier. The specific unit dosage form is dictated by and directly dependent upon (a) the unique characteristics of the particular composition and (b) the particular effect to be achieved.

20 The following non-limiting examples are provided to further illustrate the invention.

#### EXAMPLES

##### 25 I. Preparation of Synthetic FMD Vaccines.

All peptides were synthesized through the use of the Beckman 990B solid phase peptide synthesizer, using semi-automated programming. The initial functional-  
30 ization, through the first amino acid loading, and all

steps subsequent to the addition of the last amino acid, were achieved manually. All reagents unless otherwise stated were of the highest grade commercially available. Dimethylformamide (DMF) was stored over 4Å molecular  
5 sieves for one week prior to use. Dichloromethane ( $\text{CH}_2\text{Cl}_2$ ) was HPLC quality. Diisopropylethylamine (DIEA) was distilled from ninhydrin prior to use. Copoly- (styrene-1% divinylbenzene) resin, 200-400 mesh beads, was purchased from Bio-Rad Laboratories. The quality of  
10 protected amino acids, purchased from Peninsula Laboratories and Bachem, was assessed by TLC, optical rotation, amino acid analysis and mass spectral analysis prior to use.

15 A. Preparation of Aminomethyl-resin

The preparation of aminomethyl-resin was as described by A.R. Mitchell, S.B.H. Kent, M. Engelhard, R.B. Merrifield, J. Org. Chem. **43**, 2845-2852 (1978),  
20 with the exception that the desired functionalization was four-times that reported. To 100 gm of thoroughly washed resin in a three-necked round bottom flask was added 16 gm (81 mmol) of N-(hydroxymethyl)phthalimide and 2 liters of trifluoroacetic acid/methylene chloride  
25 (1:1, v/v). Under rapid overhead stirring, trifluoromethanesulfonic acid (40 ml, 0.44 mol) was slowly added. After 6 hours of reaction at 22°C, the solution was filtered and washed extensively with 4 L each of trifluoroacetic acid/methylene chloride (1:1, v/v), methylene  
30 chloride, ethanol, and finally methylene chloride.

The resin, once dried under vacuum, was refluxed for 16 hours in 2 liters of ethanol containing 5% hydrazine. The resin was filtered warm and washed with boiling ethanol (4 x 2L), methanol (4 x 2L), and methylene chloride (4 x 2L). The product was dried under vacuum to yield an amine functionalization of 0.836 mmole/gm by ninhydrin analysis [V. Sarin, S.B.H. Kent, J.P. Tam, R.B. Merrifield, Anal. Biochem. 117, 147-151 (1981)].

10 B. Preparation of Boc-Glycine-[4-(oxymethylphenyl)]  
acetic acid

The synthesis of Boc-glycine-[4-(oxymethylphenyl)]acetic acid was exactly as described by J.P. Tam, S.B.H. Kent, T.W. Wong, R.B. Merrifield, Synthesis, 955-957 (1979), with a yield of 45%.

20 C. Attachment of Derivatized Glycine to Aminomethyl Resin.

Boc-Gly-[4-(oxymethylphenyl)]acetic acid (1.4 mmol) was dissolved in 20 ml of DMF/methylene chloride (1:3, v/v) at 0°C in the presence of 1.4 mmol of 1-hydroxybenzotriazole (HOBt). An equivalent amount (1.4 mmol) of dicyclohexylcarbodiimide (DCC) was added, and the reaction mixture was stirred for 10 minutes at 0°C, after which dicyclohexylurea was removed by filtration and the supernatant added to 4 gm of aminomethyl resin suspended in 20 ml of methylene chloride. After two hours, the resin was filtered, washed three times with

40 ml of methylene chloride, and then suspended in 40 ml of 0.5M acetic anhydride/0.1M DIEA/methylene chloride. After an additional two hours, the resin was ninhydrin negative. The resin was washed three times with 40 ml  
5 of methylene chloride, and subsequently the t-butyloxy-carbonyl protecting group of glycine was removed in 40 ml of trifluoroacetic acid/methylene chloride (1:1, v/v).

10 D. Repetitive Stepwise Synthetic Protocol

In each case double coupling of each amino acid, in three-fold excess to amine, was utilized. While Asn, Gln, and Arg were coupled twice as their  
15 preformed HOBt-esters, the remaining amino acids were first coupled as their preformed symmetrical anhydrides and repeated as their preformed HOBt-esters. Neutralization, deprotection and all washing were fully automated by the following protocol, at 15 ml of  
20 solvent/gram of starting resin.

1. 5% DIEA/ $\text{CH}_2\text{Cl}_2$  (2 x 2 minutes)
2.  $\text{CH}_2\text{Cl}_2$  (6 x 1 minute)
3. First Coupling (1 x 120 minutes)
- 25 4.  $\text{CH}_2\text{Cl}_2$  (6 x 1 minute)
5. 5% DIEA/ $\text{CH}_2\text{Cl}_2$  (2 x 2 minutes)
6.  $\text{CH}_2\text{Cl}_2$  (6 x 1 minute)
7. Second Coupling (1 x 120 minutes)
8.  $\text{CH}_2\text{Cl}_2$  (6 x 1 minute)
- 30 9. 50% TFA/ $\text{CH}_2\text{Cl}_2$  (1 x 2 minutes; 1 x 20 minutes)
10.  $\text{CH}_2\text{Cl}_2$  (6 x 1 minutes)



During the course of synthesis, aliquots of the resin were removed for preparation of shortened immunogens.

5 E. Preparation of O<sub>1</sub>K, VP<sub>1</sub> (141-158)ProCysGly - -  
[21 Residue Peptide].

10 Approximately three grams of peptidyl resin representing 1.34 gm of protected O<sub>1</sub>K, VP<sub>1</sub> (141-158)-ProCysGly was treated with 30 ml of HF/p-cresol (9:1) at 0°C for 60 minutes. Upon completion of cleavage the resin was evaporated to dryness under vacuum at 0°C and subsequently washed with 5 x 50 ml of diethyl ether. The deprotected peptide was dissolved with successive  
15 acetic acid treatments: 10% HOAc (3 x 30 ml), 50% HOAc (1 x 30 ml), and finally 10% HOAc (3 x 30 ml). The peptide was freeze dried to a yield of 76%. Ellman tritration revealed a minimum free sulfhydryl content of 70%. The peptide was purified by Sephadex G-25 gel permeation chromatography in 10% HOAc following dithio-  
20 threitol (DTT) reduction to yield a highly homogeneous monomeric fraction as assessed by analytical HPLC.

25 F. Preparation of O<sub>1</sub>K, VP<sub>1</sub> [LeuProProSer(141-158)]  
ProCysGly - - [25 Residue Peptide].

30 Approximately 1.1 gm of peptidyl resin representing 0.532 gm of protected O<sub>1</sub>K, VP<sub>1</sub> [LeuProProSer(141-158)]ProCysGly was treated as in Part E to yield the desired peptide in 84% yield.

G. Preparation of O<sub>1</sub>K, VP<sub>1</sub> [(200-213)ProProSer(141-158)]  
ProCysGly - - [38 Residue Peptide].

5 Approximately 2.4 gm of peptidyl resin representing 1.29 gm of protected O<sub>1</sub>K, VP<sub>1</sub> [(200-213)ProProSer(141-158)]ProCysGly was treated as in Part E to yield the desired peptide in 97% yield.

10 H. Preparation of O<sub>1</sub>K, VP<sub>1</sub> [CysCys(200-213)ProProSer  
(141-158)]ProCysGly - - [40 Residue Peptide].

15 Approximately 2.5 gm of peptidyl resin representing 1.39 gm of protected O<sub>1</sub>K, VP<sub>1</sub> [CysCys(200-213)ProProSer(141-158)]ProCysGly was treated as in Part E to yield the desired peptide in 97% yield.

I. Preparation of O<sub>1</sub>K, VP<sub>1</sub> H-Cys-[ArgTyrAsnArgAsnAla  
(141-158)]LeuProProThr[GluAla (200-213)]OH - - [45  
Residue Peptide].

20

Protected O<sub>1</sub>K, VP<sub>1</sub> H-Cys-[ArgTryAsnArgAsnAla (141-158)]LeuProProThr[GluAla (200-213)]-peptidyl resin was treated as in Part E to produce the title peptide.

25 The amino acid analysis of these five peptides is shown in Table 1:

Table 1

## Amino Acid Composition of Synthetic FMDV Antigens

	21-Residue <sup>a</sup>	25-Residue <sup>a</sup>	38-Residue <sup>a</sup>	40-Residue <sup>b</sup>	45 Residue <sup>h</sup>
Lys	1.09(1)	1.12(1)	3.68(4)	3.74(4)	3.99(4)
His	--	--	0.91(1)	0.90(1)	1.13(1)
Arg	2.00(2)	1.99(2)	2.69(3)	2.88(3)	4.45(5)
Asp	1.08(1)	1.94(2)	2.19(2)	2.09(2) <sup>e</sup>	3.78(4)
Thr	1.05(1)	1.04(1)	2.09(2)	2.27(2) <sup>e</sup>	3.03(3)
Ser	--	0.87(1)	1.03(1)	1.18(1)	--
Glu	2.08(2)	2.13(2)	4.18(4)	4.14(4)	5.22(5)
Pro	N.D.	N.D.	N.D.	5.16(5)	4.46(4)
Gly	2.07(2)	2.12(2)	2.27(2)	2.38(2)	1.22(1)
Ala	2.14(2)	2.17(2)	3.22(3)	2.99(3)	5.12(5)
Cys	N.D.	N.D.	N.D.	2.52(3) <sup>g</sup>	N.D.
Val	2.91(3)	2.98(3)	4.83(5)	5.20(5) <sup>f</sup>	4.69(5)
Ile	--	--	0.60(1)	0.87(1)	.73(1)
Leu	2.82(3)	3.97(4)	4.34(4)	4.24(4)	5.28(5)
Tyr	--	--	--	-- .86(1)	

a. 24 hour hydrolysis in 6N HCl.

b. Average of hydrolysis at 24, 48, 72 hours in 6N HCl/DMSO.

c. Masked by cysteine oxidation product.

d. Destroyed in course of hydrolysis.

e. Determined by extrapolation to zero time of hydrolysis.

f. Determined by extrapolation to 100 hours of hydrolysis.

g. Determined as its cysteic acid derivative.

h. 21 hour hydrolysis in 6 N HCl.

I. Chemical Attachment of Peptides to KLH through SPDP.

The procedure described below for the smallest peptide studied was essentially the same for all peptides which were chemically bound to KLH. The maximum advisable loading level of SPDP [N-succinimidyl-3(2-pyridithio)propionate] to KLH was determined according to the extent of precipitation of the complex which occurred over a 60 minute reaction. It was observed that a 250 molar excess of SPDP to KLH was optimum. One-half gram of KLH ( $\sim 1 \times 10^{-4}$  mmol) was dissolved in 25 ml of 0.1M  $\text{Na}_2\text{HPO}_4$ , pH 7.2 with gentle sonication. A small quantity of insoluble material was removed by centrifugation, and to the supernatant was added 7.81 mg of SPDP ( $2.5 \times 10^{-2}$  mmol) in 625  $\mu\text{l}$  of DMF. After alkaline uptake ceased, the sample was centrifuged with the supernatant desalted by G-25 Sephadex chromatography, in 0.1M  $\text{Na}_2\text{HPO}_4$ , pH 7.2 at 20 cm/hr. The loading level achieved in the derivatization was determined to be approximately 40% by DTT reduction.

To 100 mg of the functionalized KLH ( $2 \times 10^{-6}$  mol of ligand) in 25 ml of 0.1M  $\text{Na}_2\text{HPO}_4$ , pH 7.2, was added a ten-fold excess (44 mg,  $2 \times 10^{-5}$  mol) of the (141-158)ProCysGly reduced peptide in 600  $\mu\text{l}$  of 0.1M Tris buffer at pH 7.4. The reaction was monitored by liberation of the ligand and found to be near quantitative in 60 minutes at 22°C. The sample was desalted on Sephacryl S-200 in 0.1M  $\text{NH}_4\text{HCO}_3$ , pH 7.5, and lyophilized. The yields in coupling the 21-, 25-, and 38-residue

peptides as based on limited reactant were, respectively, 84%, 85% and 68%.

J. Formulation of Peptides.

5

All peptides and peptide-KLH conjugates were dissolved in 10 mM Tris, pH 8.0, at the desired concentration and then diluted 1:1 with Freund's Complete adjuvant. The total volume as well as the antigen concentration administered to each animal is described for each experiment.

10

II. Biological Protocol

15 A. Guinea Pig Vaccination

Duncan Hartly closed colony random bred female guinea pigs, weighing 450-500 gms were used in all experiments. Each vaccine preparation was administered at the described dosage by subcutaneous injection in a volume of 0.15 ml to each member of a group of five guinea pigs. The only exception to the 0.15 ml total volume dosage was a 0.45 ml dose which was used in the antigen dose titration to deliver the highest concentration stated. The control vaccine was prepared from the O<sub>1</sub> Kaufbeuren strain, which was the one from which the published sequence data was derived. The vaccine was prepared with aluminium hydroxide gel and contained 1.66 mg of saponin per dose. Each of five guinea pigs was given one milliliter of vaccine by subcutaneous

20

25

30

injection. The challenge virus was of the O<sub>1</sub> Kaufbeuren strain as described above. It was adapted to guinea pigs by serial passage of vesicular lymph and used for challenge purposes at the fifth guinea pig passage level. Following the virus titration, a 1/20 dilution was made to give a challenge dose of 10<sup>3.5</sup> guinea pig ID<sub>50</sub>, i.e., 3,000 infective doses per animal. All test animals were given the challenge virus by intradermal inoculation of the left hind foot pad - volume 0.02 ml. Each challenged guinea pig was observed for the subsequent 7 days for development of secondary or generalized lesions on the other hind foot or hands or mouth. The criterion of protection was the absence of any secondary lesions.

#### B. Cattle Vaccination

Six to eight month old heifers of mixed breeds weighing approximately 150-180 kg which were susceptible to foot and mouth disease were used in all experiments. Each vaccine preparation was administered at the described dosage by subcutaneous injection in a volume of 3 ml to each member of a group of three cattle. In each test a group of three cattle was non-vaccinated and served as positive experimental controls. All the animals were challenged by intradermolingual injection of 10<sup>4</sup> ID<sub>50</sub> of freshly titrated O<sub>1</sub>BFS 1860 virus in a 0.1 ml dose at ten sites on the tongue. Animals were examined daily post challenge for evidence of increase in body temperature and lesion development on the tongue

and feet. Animals were deemed to be protected provided that no secondary lesions developed by seven days post-infection. All animal testing and sera evaluation was conducted at the Animal Viral Research Institute, United Kingdom.

C. Determination of Serum Neutralizing Titers (SNT)

The method employed for quantitation of serum neutralization titers was as described by Golding, S.M., Hedger, R.S., Talbot, T., Watson, J., Research in Veterinary Science 40, 142-147 (1976). All ELISA antibody titers are defined as directed against the peptide 140-160. Extremely close correlation of ELISA determinations and serum neutralization titers has been achieved in all vaccinations utilizing the described peptides.

III. Biological Results.

The most convincing method for establishing the relative potency of a series of potential synthetic peptide vaccines is to determine their minimum protective molar dose. Table 2 represents dose titration results obtained in guinea pigs with four differing immunogens. While the data is illuminating in several respects, the most important aspect is the marked enhancement in serum neutralizing titers (SNT) and protection achieved through the 38- and 40- residue peptides in comparison to that of the 21-residue

X-6060

-31-

peptide. Clearly these results indicate the unexpected advantageous properties of the compounds of this invention, which provide optimal protection upon viral challenge. The 38- and 40-residue peptides appear to be  
5 equivalent within experimental error and thereby minimize any importance of the amino terminal CysCys residues in the latter. These two peptides appear to confer complete protection at less than one-tenth to one-hundredth the dose of the 21-residue antigenic peptide.  
10 However, the unprecedented ability to protect guinea pigs against FMDV free of any stated carrier significantly simplifies the approach and eliminates a major source of variability. The most desirable synthetic agent capable of providing complete protection is that  
15 which encompasses the minimal viral structural information in a highly defined and pure state. The elimination of KLH or any like carrier while still providing protection marks a significant advance in the synthetic approach to vaccination.

20



Table 2  
Dose Titration Comparison of Synthetic FMDV in Guinea Pigs

		SNT O <sub>1</sub> K:SNT O <sub>1</sub> BFS <sup>a</sup> (No. protected: No. challenged)			
Peptide Wt <sup>b</sup>	Volume <sup>c</sup>	21-residue	21-residue-KLH	38-residue	40-residue
1. 150 nmoles	450 $\mu$ l	1/178:1/256 (5:5)	1/6:1/6 (0:5)	1/708 :1/1024 (5:5)	1/4096:1/4096 (5:5)
2. 50 nmoles	150 $\mu$ l	1/6 :1/32 (1:5)	1/6:1/6 (0:5)	1/1400:1/1400 (5:5)	1/1400:1/1400 (5:5)
3. 17 nmoles	150 $\mu$ l	1/8 :1/6 (2:5)	1/6:1/6 (0:5)	1/1024:1/1400 (5:5)	1/1400:1/2048 (5:5)
4. 6 nmoles	150 $\mu$ l	1/6 :1/6 (0:5)	1/6:1/6 (0:5)	1/708 :1/708 (4:4)	1/128 :1/90 (5:5)
5. 2 nmoles	150 $\mu$ l	1/6 :1/6 (0:5)	1/6:1/6 (0:5)	1/128 :1/256 (3:4)	1/178 :1/64 (4:5)
6. 0.6 nmoles	150 $\mu$ l	1/6 :1/6 (0:5)	1/6:1/6 (0:5)	1/11 :1/90 (5:5)	1/128 :1/178 (3:5)
7. 0.2 nmoles	150 $\mu$ l	1/6 :1/6 (0:5)	1/6:1/6 (0:5)	1/16 :1/22 (1:5)	1/64 :1/22 (1:5)

<sup>a</sup>Determined at 28 days after vaccination in pooled sera of all animals immunized with indicated peptide.

<sup>b</sup>Peptide weight for 21-residue-KLH is expressed as the amount of the synthetic peptide present in the conjugate.

<sup>c</sup>One-half of the volume consisted of Complete Freund's adjuvant.

The necessity of antigen optimization is of utmost importance, because commercialization will quite likely warrant utilization of an adjuvant of lesser potency than Complete Freund's. Table 3 reveals that at a molar dose comparable to that which provided protection with the 21-residue peptide in Complete Freund's adjuvant, the 40-residue peptide was completely protective with a commercially acceptable  $\text{Al}(\text{OH})_3$  adjuvant. The absolute weight of antigen providing full protection of guinea pigs with the latter adjuvant, in a single vaccination, is less than 1 mg.

Table 3  
Adjuvant Evaluation in Guinea Pigs with 40-Residue Peptide Vaccination

	<u>Peptide Conc</u>	<u>Adjuvant</u>	<u>No. Protected/- No. Challenged</u>
20	150 nmoles	Freund's Complete	5/5
	17 nmoles	"	5/5
	2 nmoles	"	4/5
	0.2 nmoles	"	0/5
25	150 nmoles	$\text{Al}(\text{OH})_3$	5/5
	17 nmoles	"	1/5
	2 nmoles	"	0/5
	0.2 nmoles	"	0/4

While guinea pigs have proven to be a useful model for evaluation of FMD vaccines, the ultimate test of success must be in cattle, the principal commercial target animal. Table 4 represents the results of cattle vaccination, with the 40-residue peptide, followed by the conventional challenge consisting of direct viral infection of the tongue in each vaccinated animal. The serum neutralization titers increased rapidly to reach a near maxima at Day 14, which was maintained at least through the time of challenge (Day 32). Two of the nine animals which were challenged on Day 32 were completely protected, having exhibited no signs of secondary lesions. These animals represent the first example of successful single cattle vaccination for FMD with a synthetic peptide. This attests to the potency of the immunogens of this invention. Furthermore, by serological determination of the neutralization antibody concentrations immediately prior to challenge it is not possible to predict protection. In comparison to the known minimum levels of neutralizing antibody concentrations required by a conventional FMD vaccine all of these animals should have been clearly protected. Consequently, at this time it is not acceptable to establish the efficacy of a peptide vaccine on the comparative basis of its SNT value with a conventional vaccine. Reimmunization of animals 1, 2, and 3 led to complete protection after an additional 14 days prior to challenge. Antibody titers in these animals as measured by ELISA were observed to increase approximately one half of a log unit following booster immunization. This

X-6060

-35-

40-residue peptide is less than one-fifth the size of  
any other synthetic antigen which has proven capable of  
cattle protection. Its synthetic nature will allow  
evaluation of its inherent activity in combination with  
5 various adjuvants, free of any potential low level  
bacterial potentiation.

Table 4

## Evaluation of Synthetic FMD Vaccine in Cattle

Animal No.	Peptide wt. a	SNT-Days Post Vaccination					Challenge Result <sup>b</sup>	
		0	7	14	21	26	32	
1.	1 mg	1/6	1/16	1/708	1/1024	1/1024	1/≥1400	Protected
2.	"	1/22	1/22	1/≥1400	1/≥1400	1/≥1400	1/≥1400	"
3.	"	1/16	1/45	1/1024	1/1024	1/1024	1/1024	"
4.	0.2 mg	1/11	1/22	1/≥1400	1/≥1400	1/1024	1/≥1400	Not Protected
5.	"	1/6	1/22	1/≥1400	1/≥1400	1/≥1400	1/≥1400	"
6.	"	1/32	1/32	1/≥1400	1/≥1400	1/≥1400	1/1024	"
7.	1.0 mg	1/8	1/11	1/355	1/258	1/355	1/512	"
8.	"	1/32	1/22	1/≥1400	1/≥1400	1/1024	1/1024	"
9.	"	1/11	1/22	1/1024	1/708	1/1024	1/1024	Protected
10.	5.0 mg	1/6	1/45	1/1024	1/≥1400	1/≥1400	1/≥1400	"
11.	"	1/6	1/16	1/≥1400	1/≥1400	1/≥1400	1/≥1400	Not Protected
12.	"	1/11	1/22	1/355	1/512	1/≥1400	1/1024	"
13.	None	1/6	1/16	1/11	1/6	1/90	1/32	"
14.	"	1/11	1/11	1/11	1/11	1/11	1/22	"
15.	"	1/11	1/8	1/11	1/8	1/11	1/16	"
16.	"	1/11	1/6	1/11	1/6	1/6	1/11	"

a. peptide administered was CysCys[(200-213)ProProSer(141-158)]ProCysGly as a 3 ml aqueous buffered suspension containing 50% Freund's Complete adjuvant.

b. Animals 4-16 were challenged on Day 32 following SNT determination, while the first three animals were revaccinated on Day 38 with 0.2 mg of peptide in a 3 ml suspension of Freund's Incomplete adjuvant and challenged on Day 52.

CLAIMS

1. A compound containing a sequence of  
Formula (I):

5



in which A and B are amino acid residue sequences  
comprising sequences of the foot-and-mouth disease virus  
10 VP<sub>1</sub> capsid protein serotypes, one of which contains from  
18 to 24 amino acid residues and includes the amino acid  
residue sequence in positions 141 to 158 of the O  
serotype or the equivalent sequence of other serotypes  
and the other of which contains from 14 to 20 amino acid  
15 residues and includes the amino acid residue sequence in  
positions 200 to 213 of the O serotype or the equivalent  
sequence of other serotypes;

X is H, H-Cys, or H-Cys-Cys, the sulfhydryls  
of which may be blocked or free;

20

Z is OH, Cys-OH, or Pro-Cys-Gly-OH, the  
sulfhydryls of which may be blocked or free; and

Y is a sequence of from about 2 to about 6  
amino acid residues; or a pharmaceutically-acceptable  
salt thereof.

25

2. A compound as claimed in Claim 1 in which  
X is H- or H-CysCys-.

3. A compound as claimed in Claim 1 or 2 in  
which Y is a sequence of from 2 to 4 amino acid  
residues.

30

4. A compound as claimed in Claim 3 in which  
Y contains the sequence -Pro-Pro-.

5. A compound as claimed in Claim 4 in which Y is -Pro-Pro-Ser- or Leu-Pro-Pro-Thr.

6. A compound as claimed in any one of claims 1 to 5 in which Z is -OH or -Pro-Cys-Gly-OH.

5 7.  $VP_1 (O_1K)$  [Cys-Cys(200-213)-Pro-Pro-Ser(141-158)]-ProCysGly;  $VP_1 (O_1K)$  [(200-213)-Pro-Pro-Ser-(141-158)]-ProCysGly; or a pharmaceutically acceptable salt thereof.

10 8.  $VP_1 (O_1K)$  H-Cys-[ArgTyrAsnArgAsnAla(141-158)]-LeuProProThr[GluAla(200-213)]-OH, or a pharmaceutically acceptable salt thereof.

9. A process for preparing a compound containing a sequence of Formula (I), as defined in Claim 1, which comprises

15 (a) cleaving a protecting group from a correspondingly protected compound containing the sequence of Formula (I); or

(b) culturing a microorganism transformed by an expression transfer vector containing a DNA sequence coding for a compound of Formula (I), under conditions suitable for expression of the DNA sequence coding for the compound of Formula (I), and purifying the Formula (I) compound from the lysate or culture medium of said microorganism;

25 (c) and, if desired, salifying a free compound to form a corresponding pharmaceutically-acceptable salt.

30 10. A compound containing the sequence of Formula (I), as defined in any one of claims 1 to 8, for use as a foot and mouth disease vaccine.

11. A pharmaceutical or veterinary formulation which comprises as an active ingredient a compound as defined in anyone of Claims 1 to 8 associated with one or more pharmaceutically-acceptable carriers therefor.



(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 204 480**  
**A3**

(12)

# EUROPEAN PATENT APPLICATION

(21) Application number: 86303978.0

(51) Int. Cl.<sup>3</sup>: **C 07 K 7/00**  
**C 12 P 21/02, C 12 N 15/00**  
**A 61 K 37/02**

(22) Date of filing: 27.05.86

(30) Priority: 03.06.85 US 740780

(43) Date of publication of application:  
10.12.86 Bulletin 86/50

(88) Date of deferred publication of search report: 08.02.89

(84) Designated Contracting States:  
BE CH DE FR GB IT LI LU NL SE

(71) Applicant: **ELI LILLY AND COMPANY**  
**Lilly Corporate Center**  
**Indianapolis Indiana 46285(US)**

(72) Inventor: **Dimarchi, Richard Dennis**  
**3815 Wolf Creek Circle**  
**Carmel Indiana 46032(US)**

(72) Inventor: **Brooke, Gerald Stephen**  
**2700 Dellzell Drive**  
**Indianapolis Indiana 46220(US)**

(74) Representative: **Hudson, Christopher Mark et al,**  
**Erl Wood Manor**  
**Windlesham Surrey GU20 6PH(GB)**

(54) Synthetic vaccines for foot and mouth disease.

(57) Compounds having activity in treating or preventing foot and mouth disease are disclosed. The compounds contain a sequence of the formula

X-A-Y-B-Z

in which A and B are amino acid residue sequences comprising sequences of the foot-and-mouth disease virus VP<sub>1</sub> capsid protein serotypes, one of which contains from 18 to 24 amino acid residues and includes the amino acid residue sequence in positions 141 to 158 of the O serotype or the equivalent sequence of other serotypes and the other of which contains from 14 to 20 amino acid residues and includes the amino acid residue sequence in positions 200 to 213 of the O serotype or the equivalent sequence of other serotypes;

X is H, H-Cys, or H-Cys-Cys, the sulfhydryls of which may be blocked or free;

Z is OH, Cys-OH, or Pro-Cys-Gly-OH, the sulfhydryls of which may be blocked or free; and

Y is a sequence of from about 2 to about 6 amino acid residues, or a pharmaceutically-acceptable salt thereof.

EP 0 204 480 A3



European Patent  
Office

# EUROPEAN SEARCH REPORT

0204480

Application Number

EP 86 30 3978

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Y	EP-A-0 068 693 (GENENTECH INC.) * Claims * ---	1-11	C 07 K 7/00 C 12 P 21/02
Y	EP-A-0 040 922 (BIOGEN N.V.) * Figures 9-10 * ---	1-11	A 61 K 37/02 C 12 N 15/00
Y	ENDEAVOUR, vol. 8, no. 3, 1984, pages 123-127, Pergamon Press, GB; D.J. ROWLANDS: "Synthetic peptides in foot and mouth disease vaccine research" * Page 125, right-hand column * ---	1-11	
A	JOURNAL OF IMMUNOLOGY, vol. 115, no. 6, December 1975, pages 1636-1641, The Williams & Wilkins Co., US; H.L. BACHRACH et al.: "Immune and antibody responses to an isolated capsid protein of foot-and-mouth disease virus" ---	1-11	
Y	NATURE, vol. 298, 1st July 1982, pages 30-33, Macmillan Journals Ltd; J.L. BITTLE et al.: "Protection against foot-and-mouth disease by immunization with a chemically synthesized peptide predicted from the viral nucleotide sequence" * Whole article * -----	1-11	TECHNICAL FIELDS SEARCHED (Int. Cl.4)  C 07 K 7/00 C 12 P 21/00 C 12 N 15/00 A 61 K 39/00 A 61 K 37/02
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 24-11-1988	Examiner KRASA
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document  T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			